Characterizing an Aberrant Band on Protein Electrophoresis

JANELLE CHIASERA, GEORGE RIZK, SULTAN ALATAWI

ABSTRACT

For decades, one of the most widely adopted and extensively used methods to measure serum proteins in the clinical laboratory was serum protein electrophoresis (SPEP). This method involves the separation of large molecules based on size and charge across a solid matrix (agarose gel) maintained in a vertical or horizontal plane. The resolution and surface-to-volume ratios are low in SPEP. More recently, hospital laboratories are using capillary electrophoresis (CE) for the measurement of serum proteins, mainly because of its high separation efficiency, short analysis time, low waste generation, and versatility. As opposed to SPEP, CE separates large and small molecules based on size and charge in a capillary tube. With CE, band resolution is high as well as the surface/volume ratios. Although CE has many advantages as a technique, it can lead to the development of an aberrant band appearing between the beta and gamma regions. Currently, this band is being labeled as a nonspecific band, even though the true identity of the band remains unknown. In this study, we identified 31 samples displaying an aberrant band between the beta and gamma regions on CE during a 6-month period. All samples were divided into two aliquots and frozen at -20 °C until subsequent analysis. One aliquot was treated with thrombin and reassayed using CE, and the other aliquot was used to perform Western blot. Results indicated that 23 of the 31 samples showed the presence of C-reactive protein (CRP), fibrinogen-like proteins (FGLs), or fibrinogen. Fourteen samples showed the presence of CRP, 12 samples showed the presence of FGLs, and five samples showed the likely presence of fibrinogen. The researchers concluded that the band identity is more related to acute phase proteins: CRP or FGL, or fibrinogen.

ABBREVIATIONS: CE - capillary electrophoresis, CRP - Creactive protein, FGL - fibrinogen-like protein, SPEP serum protein electrophoresis.

Clin Lab Sci 2018;31(3):163

Janelle Chiasera, The University of Alabama at Birmingham

George Rizk, The University of Alabama at Birmingham Sultan Alatawi, The University of Alabama at Birmingham Address for Correspondence: George Rizk, The University

of Alabama at Birmingham, rizk003@uab.edu